

Singularity-free Bianchi spaces with nonlinear electrodynamics.

Ricardo García-Salcedo*

CIESA-FMVZ, U. A. E. Mex., Instituto Literario 100, 50000, Toluca, México.

Nora Bretón†

Departamento de Física, Cinvestav-IPN, Apartado Postal 14-740, C.P. 07000, México, D.F., MEXICO

(Dated: February 6, 2008)

In this paper we present an analysis to determine the existence of singularities in spatially homogeneous anisotropic universes filled with nonlinear electromagnetic radiation. These spaces are conformal to Bianchi spaces admitting a three parameter group of motions G_3 . For these models we study geodesic completeness. It is shown that with nonlinear electromagnetic field some of the Bianchi spaces are geodesically complete, like G_3IX and G_3VIII ; however, completeness depends on the curvature of the space. When certain topology is assumed, Bianchi G_3IX presents geodesics that are imprisoned. It is surprising that in the linear limit (Maxwell field) the spacetimes are singularity-free even if the curvature parameter is zero.

PACS numbers: 98.80.Hw, 11.10.Lm, 04.20.Dw, 04.20.Jb

I. INTRODUCTION

General Relativity breaks down at certain points called singularities. The singularity theorems, that were shown for the first time 30 years ago by Stephen Hawking, Robert Geroch and Roger Penrose [1], gave the bad news that under certain conditions the singularities of the spacetime would appear in the future or in the past. An important subject that is not yet determined is the nature of the singularities of the spacetime in general relativity.

There are several criteria that tell us if a spacetime is singular, such as energy conditions or the divergence of the curvature invariants. However, these conditions are only necessary but not sufficient. If one of them fails to hold, one is ignorant if the spacetime possesses a singularity or not. The common characteristic of singular spacetimes is the existence of incomplete causal geodesics. We shall adopt as definition of the absence of singularities the completeness of causal geodesics (g-completeness), i. e. no observer in free fall leaves the spacetime in a finite proper time. We note, however, that not even the geodesic completeness guarantees the absence of singularities (see a counterexample by Geroch [2], of a geodesically complete spacetime with incomplete non-geodesic curves).

Mars and Senovilla presented a Bianchi IX model with an energy-momentum tensor of perfect fluid type [3] that is completely regular everywhere in the energy density and pressure and is geodesically complete. This model provides an example of how the presence of matter can help to avoid the initial singularity while keeping physical compelling dominant energy conditions.

Recently, more studies about singularity-free Bianchi

type IX models have arisen; for example, Berger used the method of consistent potentials [4] to explain how minimally coupled (classical) scalar field can suppress mixmaster oscillations [5] when approaching the singularity of Bianchi IX cosmology. Toporensky and Ustiansky studied the possibility of non-singular transition from contraction to expansion in the dynamics of the Bianchi IX cosmological model with minimally coupled massive real scalar field [6].

In this paper we study singularities in spatially homogeneous but anisotropic spaces where the energy-momentum tensor corresponds to (source-free) nonlinear electromagnetic field, in particular, Born-Infeld field. For these spaces the energy momentum tensor satisfies the dominant energy condition and the matter does not diverge; three cases possess regular Weyl invariants. These conditions point to the absence of singularities provided the spaces do pass the test of geodesic completeness. The assumed spacetime includes particular cases of Bianchi spaces. The analysis of the geodesics lead us to conclude that some of the studied Bianchi spaces avoid the singularity, like Bianchi G_3IX and G_3VIII , while Bianchi G_3II might present singularity. Spaces like Bianchi G_3III and G_3I as well as a particular case of Kantowski-Sachs present singularity in the curvature, i. e. the invariants diverge after a finite time, however, in two cases the geodesics are complete.

The solutions investigated in this papers have already been discussed. Approximate solution was presented in [7] and the possibility of (NLED) as source of inflation was addressed in [8]. The singularity structure is studied in the present paper.

Before getting into the main subject, we give a summary of non-linear electrodynamics (NLED) in Sec. II. In Sec. III the studied solutions of the coupled Einstein-Born-Infeld (EBI) equations are presented along with comments on their geometric structure, energy conditions and scalar invariants. In Sec. IV we analyze the geodesics and the nature of the singularities that appear

*Electronic address: rgs@uaemex.mx

†Electronic address: nora@fis.cinvestav.mx

in the models. In Sec. V we address the limit of linear electrodynamics (Einstein-Maxwell) and it turns out that for this field all solutions are geodesically complete. In the final section we draw some conclusions.

II. BORN-INFELD NONLINEAR ELECTRODYNAMICS

One of the motivations to explore spacetimes with nonlinear electromagnetic fields is that in early epochs of the universe the magnetic fields exceeded some 10^{15}G . At such values of the electromagnetic field, the interaction of photons with themselves becomes important and classical electrodynamics is not valid anymore [9]. To overcome the use of quantum electrodynamics (QED) we shall employ the Born-Infeld theory.

The nonlinear electrodynamic theory presented by Born and Infeld (1934) possesses a Lagrangian which takes into account quantum electrodynamical corrections, this fact was noted by Euler and Heisenberg soon after the proposal by Born and Infeld and later on it was remarked by Schwinger. In the papers by Heisenberg and Euler [10] and by Drummond and Hathrell [11], it was demonstrated that the Born-Infeld lagrangian has the same dependence in the electromagnetic invariants than QED for the one-loop approach. On the other side, based on very general considerations, Schwinger [12] obtained an expression for the lagrangian that coincides, within numerical factors of order unity, with the BI lagrangian in the same approximation. Therefore, EBI theory should give us, at least qualitatively, characteristics of early universes that could not be obtained within classical electrodynamics.

Moreover, lately the BI type action has arisen in string theory: it happens that open strings with Dirichlet boundary conditions (D-branes) are described by an effective action of Dirac-Born-Infeld, that is related to the effective action of BI type in the theory of open strings. The action to low energies corresponding to these objects, called branes, can be written like a term of BI plus a term of the Weiss-Zumino or Chern-Simmons type [13]. The BI strings are solitons that represent terminal strings on branes.

On the other side, in relation with cosmological models with electromagnetic fields, Brill [14] studied Einstein-Maxwell solutions in a homogeneous and nonisotropic universe. His solution is a generalization of Taub's vacuum universe and represents a closed universe of topology $\mathbb{R} \times S^3$ filled with gravitational and electromagnetic radiation. This universe expands in two spatial directions and contracts along the third one; moreover its invariants are finite. Also Hughston and Jacobs [15] investigated the behavior of source-free, homogeneous, electromagnetic fields in Bianchi type cosmologies. They showed that there can be no pure electric (magnetic) fields in types VIII, IX, V, VI ($h \neq 0$) and VI ($h \neq -1$). Types III, VI ($h = -1, h = 0$) allow one independent component of the

magnetic field. Type II allows two independent magnetic field components while type I puts no constraints on the magnetic field components. Note that results in [15] were obtained for Maxwell fields in the Bianchi geometry, non coupled to Einstein equations. In a previous paper, Hacyan [16] investigated the effect of BI electrodynamics in the initial singularity of a Bianchi I space. It turned out in that case that the presence of a BI field smoothed the oscillations of the metric functions when approaching the singularity. In the cases addressed in the present paper, Bianchi types $G_3\text{I}$, $G_3\text{III}$ as well as Kantowski-Sachs admit an electric or magnetic non linear field (not both) and $G_3\text{II}$, $G_3\text{VIII}$ and $G_3\text{IX}$ admit both components, moreover, the presence of the electromagnetic field prevents the occurrence of singularities in some Bianchi spaces.

A. The Born-Infeld field parameterization

We describe in what follows the main characteristics of the BI formalism used to determine the EBI solutions.

The Born-Infeld (BI) nonlinear electromagnetic theory is self-consistent and satisfies all natural requirements. Its lagrangian depends on the BI constant b , the maximum field strength; it also depends in nonlinear form on the electromagnetic invariants, P and Q , of the antisymmetric tensor $P^{\mu\nu}$, the generalization of the electromagnetic tensor $F^{\mu\nu}$; the BI lagrangian is given by

$$L_{BI} = -\frac{1}{2}P^{\mu\nu}F_{\mu\nu} + \mathcal{H}(P, Q) \quad (1)$$

where $\mathcal{H}(P, Q)$ is the so called structural function. The structural function is constrained to satisfy some physical requirements: (i) the correspondence with the linear theory of Maxwell in the limit when b goes to infinity ($\mathcal{H}(P, Q) = P + O(P^2, Q^2)$); (ii) the parity conservation ($\mathcal{H}(P, Q) = \mathcal{H}(P, -Q)$); (iii) the positive definiteness of the energy density ($\mathcal{H}_{,P} > 0$) and, (iv) the requirement of the timelike nature of the energy flux vector ($P\mathcal{H}_{,P} + Q\mathcal{H}_{,Q} - \mathcal{H} \leq 0$). Conditions (iii) and (iv) amount to the fulfilment of the dominant energy condition (DEC) [1]: $T_{\mu\nu}u^\mu u^\nu > 0$, for every time direction u_μ , ($u^\mu u_\mu < 0$) and $T_{\mu\nu}u^\mu$ is a nonspacelike vector. The energy-momentum tensor in terms of the structural function is given by:

$$T_{\mu\nu} = -\mathcal{H}_{,P} P_{\mu\alpha} P^\alpha{}_\nu + g_{\mu\nu} (2P\mathcal{H}_{,P} + Q\mathcal{H}_{,Q} - \mathcal{H}). \quad (2)$$

The structural function for the BI field is given by

$$\mathcal{H}(P, Q) = b^2 - \sqrt{b^4 - 2b^2 P + Q^2}, \quad (3)$$

In the linear limit, which is obtained by taking $b \rightarrow \infty$, then $\mathcal{H} = P$ and $P_{\mu\nu} = F_{\mu\nu}$. $P_{\mu\nu}$ and $F_{\mu\nu}$ are related through the material or constitutive equations:

$$F_{\mu\nu} = \mathcal{H}_{,P} P_{\mu\nu} + \mathcal{H}_{,Q} \check{P}_{\mu\nu}, \quad (4)$$

where $\check{P}_{\mu\nu} = -\frac{1}{2}\epsilon_{\mu\nu\alpha\beta}P^{\alpha\beta}$ is the dual of $P_{\mu\nu}$. An extensive treatment on nonlinear electrodynamics in curved spacetimes can be consulted in [17].

In the null tetrad formalism [18], and dealing with a Petrov type-D metric, one can always align the directions of the real null vectors e^3 and e^4 along the double Debever-Penrose (DP) vectors. We shall assume that the eigenvectors of F_{ab} (and consequently the ones of P_{ab}) are also aligned in the direction of the DP vectors. Hence the nonvanishing components of F_{ab} are F_{12} and F_{34} (P_{12} and P_{34}). We shall adopt the following parameterization:

$$P_{12} = iH, \quad P_{34} = D, \quad (5)$$

where D is the electric displacement and H is the magnetic field, the invariants acquire the form:

$$P = \frac{P^{ab}P_{ab}}{4} = -\frac{D^2 - H^2}{2}, \quad Q = \frac{P^{ab}\check{P}_{ab}}{4} = iHD, \quad (6)$$

In this parameterization the structural function turns out to be

$$\begin{aligned} \mathcal{H} &= b^2 - \sqrt{b^4 + b^2(D^2 - H^2) - H^2D^2} \\ &= b^2 - \sqrt{(b^2 + D^2)(b^2 - H^2)}. \end{aligned} \quad (7)$$

The parameterization (5) has the advantage that there are only two nonvanishing components of the energy momentum tensor:

$$\begin{aligned} T_{34} &= 2b^2 \left[\frac{b^2 + D^2}{\sqrt{(b^2 + D^2)(b^2 - H^2)}} - 1 \right] = 2b^2(e^{-\nu} - 1), \\ T_{12} &= 2b^2 \left[\frac{b^2 - H^2}{\sqrt{(b^2 + D^2)(b^2 - H^2)}} - 1 \right] = 2b^2(e^{\nu} - 1) \end{aligned} \quad (8)$$

where we have defined:

$$e^{\nu} = \sqrt{\frac{b^2 - H^2}{b^2 + D^2}} \quad (9)$$

The condition that e^{ν} be real imposes the restriction that $H < b$, i. e. the fields do not reach the maximum allowed field b .

III. SPATIALLY HOMOGENEOUS SPACETIMES WITH NON-LINEAR ELECTROMAGNETIC FIELD

Since the energy momentum tensor has two nonvanishing eigenvalues, Eqs. (8), may be coupled to spacetimes with two preferred directions, i. e. spacetimes of type D in the classification by Petrov. We have found solutions to the coupled Einstein-Born-Infeld equations for spatially homogeneous and anisotropic spaces, among them there are several cases of Bianchi metrics.

TABLE I: Values of $h(z)$ and ϵ for which the studied metric is conformal to the indicated Bianchi types

σ^1	$h(\theta) = 1 - \epsilon z^2(\theta)$	ϵ	Bianchi type
dX	$\sin^2 \theta$	1	K-S
dX	1	0	G ₃ I
dX	$\sinh^2 \theta$	-1	G ₃ III
$dX + 2l \cos \theta dy$	$\sin^2 \theta$	1	G ₃ IX
$dX + 2l z dy$	1	0	G ₃ II
$dX + 2l \cosh \theta dy$	$\sinh^2 \theta$	-1	G ₃ VIII

The studied line element is of the form

$$\phi(t)^2 ds^2 = -\frac{dt^2}{s(t)} + s(t)(dx + 2l z dy)^2 + \frac{dz^2}{h(z)} + h(z)dy^2, \quad (10)$$

where x and y are ignorable coordinates, $h = 1 - \epsilon z^2$, with ϵ and l being constants; $s = s(t)$ and $\phi = \phi(t)$ are the metric functions that are determined from the Einstein-Born-Infeld coupled equations. Chosing the null tetrad in the form:

$$\begin{aligned} e^1 &= \bar{e}^2 = \frac{1}{\sqrt{2}\phi} \left[\frac{dz}{\sqrt{h}} + i\sqrt{h}dy \right], \\ e^3 &= \frac{1}{\sqrt{2}\phi} \left[\frac{dt}{\sqrt{s}} + \sqrt{s}(dx + 2l z dy) \right], \\ e^4 &= \frac{1}{\sqrt{2}\phi} \left[-\frac{dt}{\sqrt{s}} + \sqrt{s}(dx + 2l z dy) \right], \end{aligned} \quad (11)$$

The metric function $s(t)$ turns out to be

$$s(t) = \phi^3 \dot{\phi} \left[C_1 + 2b^2 \int \frac{(1 - e^{\nu})}{\dot{\phi}^2 \phi^4} dt + \epsilon \int \frac{dt}{\dot{\phi}^2 \phi^2} \right], \quad (12)$$

where dot simbolizes derivative with respect to time t , C_1 is an integration constant related to the vacuum case, b is the BI parameter and ϵ is the constant curvature parameter. Moreover, $\phi(t)$ must satisfy the differential equation $\ddot{\phi} + l^2 \phi = 0$.

The system of EBI equations admits two possible solutions for the electromagnetic function $\nu(t)$, Eq. (9),

$$e^{\nu(t)} = \sqrt{1 \pm \phi^4(t)}. \quad (13)$$

Both solutions present regions in which $s(t) > 0$ passing then to the opposite sign, $s(t) < 0$. The regions with $s(t) > 0$ admit a cosmological interpretation as spatially homogeneous spaces. For $s(t) < 0$ the signature of (10) changes corresponding to stationary spaces (t direction changes from timelike to spacelike). We shall study the case $e^{\nu} = \sqrt{1 - \phi^4}$. The condition e^{ν} to be real imposes the restriction that $\phi^4 < 1$ and it determines

the range of the coordinate t . The electromagnetic fields are homogeneous and constant on spacelike hypersurfaces $\{t = \text{const}\}$:

$$\begin{aligned} D(t) &= \phi^2(t) \cos \left[2l \int \frac{dt}{\sqrt{1-\phi^4}} \right], \\ B(t) &= -\phi^2(t) \sin \left[2l \int \frac{dt}{\sqrt{1-\phi^4}} \right], \end{aligned} \quad (14)$$

or in terms of the electromagnetic tensor:

$$\begin{aligned} \phi^2(t) F_{zy} &= B(t) = e^\nu H(t), \\ \phi^2(t) F_{tx} &= E(t) = e^{-\nu} D(t), \end{aligned} \quad (15)$$

The results in subsections III.B and III.C were derived in [7] but are included here for completeness.

The line element (10) can be written in terms of a synchronous time coordinate if we transform $\frac{dt^2}{s(t)} = dT^2$ and with $\sigma^1 = dx + 2l z dy$:

$$dS^2 = -dT^2 + s(\sigma^1)^2 + \frac{dz^2}{h} + h dy^2, \quad (16)$$

Written in the form (16) the Bianchi types can be identified according to Table I (cf. Eqs. (13.1)-(13.2) in [19]). In case Bianchi G₃IX the metric (16) is a particular case of Taub's universe (cf. Eqs. (6.29) or (8.5) in [20]).

A. Bianchi cases included

The spacetime (10) possesses four Killing vectors. The four generator group G_4 has invariant subgroups G_3 that can be spatial rotations or translations (S_3 or T_3). In particular, we shall pay attention to the hypersurface-homogeneous spacetimes that can be identified as Bianchi spaces I, II, III, VIII and IX and one of them as Kantowski-Sachs.

The Killing vectors X_a are classified according to the values of the curvature parameter, ϵ that can be 1, 0, -1:

i) $\epsilon = 0$, $h = 1$

$$\begin{aligned} X_1 &= \partial_y, \quad X_2 = \partial_x, \\ X_3 &= -\partial_z/2l + y\partial_x, \\ X_4 &= z\partial_y - y\partial_z + l(y^2 - z^2)\partial_x. \end{aligned} \quad (17)$$

ii) $\epsilon = -1$, $h = 1 + z^2$,

$$\begin{aligned} X_1 &= \partial_y, \quad X_2 = \partial_x, \\ X_3 &= \sqrt{h} \cosh y \partial_z - \frac{\sinh y}{\sqrt{h}} (2ly\partial_x + z\partial_y), \\ X_4 &= \sqrt{h} \sinh y \partial_z - \frac{\cosh y}{\sqrt{h}} (2ly\partial_x + z\partial_y), \end{aligned} \quad (18)$$

iii) $\epsilon = 1$, $h = 1 - z^2$,

$$\begin{aligned} X_1 &= \partial_y, \quad X_2 = \partial_x, \\ X_3 &= \sqrt{h} \cos y \partial_z - \frac{\sin y}{\sqrt{h}} (2ly\partial_x - z\partial_y), \\ X_4 &= \sqrt{h} \sin y \partial_z + \frac{\cos y}{\sqrt{h}} (2ly\partial_x - z\partial_y) \end{aligned} \quad (19)$$

Special cases of Bianchi metrics with a particular topology can be obtained as follows:

i) $\epsilon = 0$

In the metric (10) with $\epsilon = 0$ we obtain $h(z) = 1$ and the line element is

$$\phi(t)^2 ds^2 = -\frac{dt^2}{s(t)} + s(t)(dx + 2lz dy)^2 + dz^2 + dy^2, \quad (20)$$

where the ranges of all the coordinates are from $-\infty$ to ∞ , this is a non-compact spacetime with topology R^4 that corresponds to Bianchi II. If $l = 0$, the metric becomes diagonal and can be identified as a Bianchi I,

$$\phi(t)^2 ds^2 = -\frac{dt^2}{s(t)} + s(t)dx^2 + dz^2 + dy^2. \quad (21)$$

ii) $\epsilon = -1$

In the metric (10) we transform $z \rightarrow \sinh \theta$ obtaining $h(\theta) = \cosh^2 \theta$ and the line element becomes

$$\begin{aligned} \phi(t)^2 ds^2 &= -\frac{dt^2}{s(t)} + s(t)(dx + 2l \sinh \theta dy)^2 + \\ &\quad d\theta^2 + \cosh^2 \theta dy^2, \end{aligned} \quad (22)$$

where the ranges of all the coordinates are from $-\infty$ to ∞ , this is a non-compact spacetime with topology R^4 that corresponds to Bianchi VIII. If $l = 0$ in the metric (10) and we transform $z \rightarrow \cosh \theta$, $h(\theta) = \sinh^2 \theta$, then the line element is

$$\phi(t)^2 ds^2 = -\frac{dt^2}{s(t)} + s(t)dx^2 + d\theta^2 + \sinh^2 \theta dy^2, \quad (23)$$

that can be identified as a Bianchi III space.

iii) $\epsilon = 1$

In the metric (10) we transform $z \rightarrow \sin \theta$ and we obtain $h(\theta) = \cos^2 \theta$ and the line element as

$$\phi(t)^2 ds^2 = -\frac{dt^2}{s(t)} + s(t)(dx + 2l \sin \theta dy)^2 + d\theta^2 + \cos^2 \theta dy^2, \quad (24)$$

where the ranges of the coordinates are $-\infty < t < \infty$, $0 \leq \theta \leq \pi$; if $0 \leq y \leq 2\pi$, $0 \leq x \leq 4\pi$ it is a compact spacetime with topology $R \times S^3$ that corresponds to

Bianchi IX. For $l = 0$ in metric (10) and changing coordinates to $z \rightarrow \cos \theta$ and with $\epsilon = 1$ we obtain $h(\theta) = \sin^2 \theta$ and the line element transforms into

$$\phi(t)^2 ds^2 = -\frac{dt^2}{s(t)} + s(t)dx^2 + d\theta^2 + \sin^2 \theta dy^2, \quad (25)$$

it is a spatially-homogeneous solution which admits no simply-transitive G_3 that can be identified as a Kantowski-Sachs space.

B. Energy conditions

According to Raychaudhuri, the equation that governs the volume expansion of geodesic congruences, θ , is

$$\frac{d\theta}{d\lambda} = -R_{ab}V^aV^b + 2\omega^2 - 2\sigma^2 - \frac{\theta^2}{3} + \dot{V}^\alpha_{;\alpha} \quad (26)$$

where V^α is a tangent vector to the geodesics, R_{ab} is the Ricci tensor, ω is the vorticity, σ is the shear and $\dot{V}^\alpha$ is the acceleration of geodesics (or flow lines in a fluid); we shall consider that $\omega = 0$.

From Eq. (26) it can be seen that the expansion θ of a timelike geodesic congruence with zero vorticity will monotonically decrease along a geodesic if, for any timelike vector V^a , $R_{ab}V^aV^b \geq 0$. This is the strong energy condition (SEC), as stated in [21], it says: “A spacetime satisfies the strong energy condition if $R_{ab}V^aV^b \geq 0$, for all causal vectors V^a ”. The term involving the Ricci tensor, R_{ab} , in Eq. (26) induces contraction of the geodesic lines, indicating that the focusing of neighbouring geodesics requires SEC.

For our Bianchi spaces the two nonvanishing components of the traceless Ricci tensor, in terms of the matter content, are

$$R_{12} = 2b^2(1 - e^{-\nu}), \quad R_{34} = 2b^2(1 - e^{-\nu}). \quad (27)$$

If we consider the congruence $V^a = \phi\sqrt{s}\partial_t$, or in the null tetrad formalism, $V^3 = V_4 = \frac{1}{\sqrt{2}}$, $V^1 = V^2 = 0$, for the BI field SEC amounts to $R_{ab}V^aV^b = R_{34} = 2b^2(1 - e^{-\nu}) < 0$, that clearly violates SEC.

The violation of SEC is not exceptional, we remind that minimally coupled scalar field violates SEC and indeed curvature-coupled scalar field theory also violates SEC. This point has been discussed recently in [21] and [22]. The aspect of interest for us is that the violation of this condition could imply the absence of an initial singularity since the fulfilling of SEC is related with the convergence of neighbouring geodesics.

On the other side, the dominant energy condition (DEC) that states “for every timelike vector, V^a , $T_{ab}V^aV^b \geq 0$ and $T_{ab}V^a$ is a nonspacelike vector” [1], is indeed fulfilled by the BI energy-momentum tensor,

Eqs. (2)-(3). DEC means that to any observer the local energy density appears nonnegative and the local energy flow vector is non-spacelike. In the BI case DEC is a requirement of a physically reasonable nonlinear structural function $\mathcal{H}$ and its fulfilment is guaranteed by conditions (iii) and (iv) as stated in Sec. II.A.

C. Scalar Polynomial Invariants

For these solutions the only nonvanishing Weyl scalar is given by

$$\Psi_2 = \frac{\phi^2}{12} (\ddot{s} - 8sl^2 + 2\epsilon + 6il\dot{s}). \quad (28)$$

The scalar polynomial invariants constructed from Ψ_2 are given by $I = 3\Psi_2^2$ and $J = -\Psi_2^3$, in such a manner that if Ψ_2 is unbounded then both I and J must be unbounded and a curvature singularity occurs. The regularity and boundedness of Ψ_2 depends on the proper for $\ddot{s}(t)$, $\dot{s}(t)$ and $s(t)$. In case of a suspected singular point, we must check if Ψ_2 is regular when evaluated at the suspicious point. If Ψ_2 remains bounded then it proceeds to investigate if the geodesics are complete.

One could think that regularity of the curvature invariants might be helpful to avoid singularities but there are known examples of spacetimes with regular invariants, such as Taub-NUT, that enclose geodesics that are not complete in their affine parameterization due to the phenomenon called *imprisoned incompleteness* [1], in which a non-spacelike curve as it follows to its future enters and remains within a compact set. In spite that Hawking himself asserted that “the kind of behavior which occurs in Taub-NUT space cannot happen if there is some matter present”, when studying a spacetime with regular invariants we must show that in fact geodesics are complete if one wants to assert that the space is singularity-free.

D. Linear limit

The linear (Maxwell) limit can be obtained from the solution Eq. (12) taking the limit $b \rightarrow \infty$, obtaining

$$s(t) = \phi^3 \dot{\phi} [C_1 + C^2 \int \frac{dt}{\dot{\phi}^2 \phi^4} + \epsilon \int \frac{dt}{\dot{\phi}^2 \phi^2}], \quad (29)$$

where $C^2 = H^2 + D^2$ is related to the energy density of the field; in this case the electromagnetic field is given by

$$H = C \sin(2lat), \quad D = C \cos(2lat), \quad (30)$$

where a is a constant. In Fig. 1 are displayed both metric functions, the linear (Maxwell) and Born-Infeld, the values of the constants correspond to a case of Bianchi G_3 IX. In general the BI field smooths oscillations of




FIG. 1: The metric functions $s(t)$ when the BI field is present (continuous curve) and in the linear limit (Maxwell field) (dashed curve) are shown. In general, the presence of BI field smooths oscillations of the metric function. The plot corresponds to Bianchi G_3 IX, the values of the constants are $l = 1$, $A = 0.8$, $C_1 = 0.5$, $b = 0.85$, $\epsilon = 1$ and the value of the constant linear field is $C = 0.85$.

the metric function $s(t)$. It is surprising that with linear electromagnetic field as matter the spacetimes are singularity-free. We shall address geodesic completeness for this case in Sec. V.

IV. ANALYSIS OF SINGULARITY STRUCTURE

Hawking and Ellis [1] discussed Bianchi I spaces and gave a theorem asserting that singularities will occur in all non-empty spatially homogeneous models in which the timelike convergence condition ($R_{ab}u^a u^b \geq 0$) is satisfied. However, as we showed in section III.B, in the studied case the timelike convergence condition is not fulfilled. Therefore the possibility of the absence of singularity persists.

For our Bianchi spaces with nonlinear electrodynamics there are two branches characterized by the value of the constant l that appears in the non-diagonal term of the metric, $(dx + 2l z dy)$. In the case $l = 0$ the conformal function $\phi(t)$ fulfills the equation $\ddot{\phi} = 0$, then $\phi = At + B$

where A and B are constants. In the case $l \neq 0$ the equation for $\phi(t)$ is $\ddot{\phi} + l^2 \phi = 0$, with solutions $\phi = A \cos(lt + B)$, A and B being constants. The analytical expression of the metric function $s(t)$ depends on the conformal function $\phi(t)$ and its behavior is different in each case.

We will consider a spacetime as singularity-free when it is geodesically causal complete (g-completeness), that means intuitively that an observer in free falling does not leave the spacetime in a finite proper time; equivalently, that every geodesic can be extended to arbitrary values of its affine parameter. The demonstration of geodesic completeness is not simple in most cases [21]. Nevertheless, the analysis of the geodesic extension can be simplified when there exist constants of motion associated to Killing vectors.

The method that we use to analyze the completeness is to obtain the first order equations for the derivatives of the coordinates with respect to an affine parameter, x, τ , t, τ , y, τ , z, τ , using the constants of motion. Then showing that these first derivatives are bounded, this implies that the corresponding geodesic curves $x(\tau)$, $t(\tau)$, $y(\tau)$, $z(\tau)$, are complete [23].

We shall analyze geodesic completeness of the spacetimes in the following subsections, first for the class of non-diagonal metrics, $l \neq 0$ that includes Bianchi G_3 II, G_3 VIII and G_3 IX and then in the next subsection for the diagonal metrics Kantowski-Sachs and Bianchi spaces G_3 I and G_3 III.

A. Bianchi G_3 II, G_3 VIII and G_3 IX

In the case that $l \neq 0$ the equation for ϕ is $\ddot{\phi} + l^2 \phi = 0$, with solution $\phi = A \cos(lt + B)$, A, B being constants; the sector (t, x) of the metric is non diagonal and the cases included are Bianchi G_3 VIII and G_3 IX for $\epsilon = -1, 1$, respectively and G_3 II when $\epsilon = 0$.

The metric function $s(t)$, Eq. (12) can be integrated in exact form; with $\phi = A \cos t$ (taking $l = 1$ and $B = 0$) we obtain a cumbersome expression in terms of $\sin t$, $\cos t$ and elliptic functions,

$$\begin{aligned}
 s(t) &= -A^4 C_1 \cos^3 t \sin t + \epsilon (\cos^4 t - \cos^2 t \sin^2 t) + \\
 &\quad \frac{2b^2}{A^2} \left\{ \cos^4 t - \frac{5}{3} \cos^2 t \sin^2 t - \frac{1}{3} \sin^2 t - \frac{1}{6} (2 \cos 2t + \cos 4t) \sqrt{(2 - A^2 - A^2 \cos^2 t)(2 + A^2 + A^2 \cos^2 t)} + \right. \\
 &\quad \left. \frac{8}{3} A^2 \cos^4 t \sin^2 t \sqrt{\frac{(2 + A^2 + A^2 \cos^2 t)}{(2 - A^2 - A^2 \cos^2 t)}} - \frac{2}{3} \sin t \cos^3 t [\sqrt{1 + A^2} E[\alpha, \beta] - \left(\frac{4 - A^4}{\sqrt{1 - A^2}} \right) F[\gamma, \beta]] \right\}, \\
 \alpha &= \arcsin \left(\frac{\sqrt{2} \sin t}{\sqrt{2 - A^2 - A^2 \cos 2t}} \right), \quad \beta = \frac{2A^2}{1 + A^2}, \quad \gamma = \operatorname{arcsinh} \left(\frac{(1 - A^2) \sqrt{2 + A^2 + A^2 \cos 2t}}{2 \sin t} \right),
 \end{aligned} \tag{31}$$

E and F are the elliptic integrals of second and first

kind, respectively. From the expression (31), it is clear




FIG. 2: This is the characteristic plot for the metric function $s(t)$ and its derivatives, $\dot{s}(t), \ddot{s}(t)$ in the case $l \neq 0$. The three functions are periodic and bounded. The corresponding Weyl scalar, Ψ_2 does not diverge in the full range $-\infty < t < \infty$. In this graphic the values of the constants are $A = 0.8, C_1 = 0.5, b = 0.85$ and $\epsilon = 1$. For $\epsilon = 0, -1$ the behavior is qualitatively the same.

that $s(t)$ is a periodic and bounded function. There are values of t for which the metric function $s(t)$ is null and there the metric (10) is singular. For $\phi(t) = A \cos(t)$, these values are $t = \pm(2n+1)\pi/2, n = 0, 1, 2, \dots$ and there we might have a singularity. There are additional zeroes when negative terms cancel the positive ones, however, the position of these zeroes depends on the balance between the distinct parameters, $C_1, b^2/A^2, A^2$ and ϵ .

The characteristic behavior of the functions $s(t), \dot{s}(t), \ddot{s}(t)$ is illustrated in Fig.2. The three functions $s(t), \dot{s}(t), \ddot{s}(t)$ are periodic and bounded, that do not diverge over all the range $-\infty < t < \infty$. As a consequence, the Weyl scalar Ψ_2 , Eq. (28), is also finite over all the range $-\infty < t < \infty$ and so are its derived invariants. To dilucidate the behavior of the spacetime at $t = \pi/2$ we shall investigate the completeness of geodesics.

For each Killing vector X_i there exists a conserved quantity $x_{,\tau}^\mu X_{\mu} = x_{,\tau}^\mu g_{\mu\nu} X^\nu = \text{const}$, where $x^\mu(\tau)$ is a geodesic parametrized by τ . Since ∂_x and ∂_y are Killing vectors, it implies that there are constants K_1, K_2 given by

$$\begin{aligned} K_1 &= g_{xx}x_{,\tau} + g_{xy}y_{,\tau}, \\ K_2 &= g_{yy}y_{,\tau} + g_{yx}x_{,\tau}, \end{aligned} \quad (32)$$

From Eqs. (32) we determine $x_{,\tau}$ and $y_{,\tau}$:

$$\begin{aligned} y_{,\tau} &= \frac{\phi^2}{h}(K_2 - MK_1), \\ x_{,\tau} &= \frac{\phi^2 K_1}{s} - My_{,\tau}, \end{aligned} \quad (33)$$

where $M = 2lz$. From Eqs. (33) we can analyze completeness for $x(\tau)$ and $y(\tau)$. All the coefficients in Eqs. (33) are finite ($h = 1 - \epsilon z$) except for the first term in

$x_{,\tau}$, since there will be divergences if s/ϕ^2 is null. We shall analyze this term with detail.

Taking the limit $t \rightarrow \pi/2$,

$$\lim_{t \rightarrow \pi/2} \left(\frac{s}{\phi^2} \right) = -\epsilon, \quad (34)$$

therefore, only in the case that $\epsilon = 0$, the geodesics $x(\tau)$ can be incomplete. This case corresponds to spaces Bianchi II. Even in this case we can have complete geodesics when the constant $K_1 = 0$, that means,

$$K_1 = \frac{s}{\phi^2}(x_{,\tau} + My_{,\tau}) = 0, \quad (35)$$

in that case the geodesics $x_{,\tau} = -My_{,\tau}$, but the equation for $y_{,\tau}$ is given in terms of bounded coefficients, also $M = 2lz$ is bounded, then in this particular case of $\epsilon = 0$, the geodesics $x(\tau)$ are complete. However the generic behavior when $\epsilon = 0$ is geodesically incomplete.

To obtain the equations for $t_{,\tau}$ and $z_{,\tau}$ we shall use the rest of the Killing vectors, quoted in Sec. III.A. These Killing vectors are different depending on the value of the curvature parameter ϵ . We separate in three cases: $\epsilon = 1, -1, 0$.

For $\epsilon = 1$ the Killing vectors Eqs. (19) imply the existence of two conserved quantities, K_3 and K_4 that permit determine $z_{,\tau}$ as

$$z_{,\tau}^2 + \phi^4(zK_2 - 2lK_1)^2 - \phi^4h(K_3^2 + K_4^2) = 0, \quad (36)$$

On the other hand, we can also use the line element, to obtain $t_{,\tau}$,

$$ds^2 = \delta = \frac{z_{,\tau}^2}{\phi^2 h} + \frac{h}{\phi^2} y_{,\tau}^2 - \frac{t_{,\tau}^2}{\phi^2 s} + \frac{s}{\phi^2} (x_{,\tau} + My_{,\tau})^2, \quad (37)$$

where δ is 1 for spacelike geodesics, 0 for null and -1 if the geodesics are timelike.

Substituting the constants of motion we have,

$$\begin{aligned} t_{,\tau}^2 &= s\phi^4\{(K_3^2 + K_4^2) - (zK_2 - 2lK_1)^2/h + \\ &\quad (K_2 - K_1M)^2/h\} + \phi^4K_1^2 - \phi^2s\delta. \end{aligned} \quad (38)$$

In the equations for $t_{,\tau}^2$ and $z_{,\tau}^2$ all the coefficients are bounded, then the geodesic curves $t(\tau)$ and $z(\tau)$ are complete.

In an analogous way can be determined $t_{,\tau}$ and $z_{,\tau}$ for the case $\epsilon = -1$. Using the Killing vectors X_3 and X_4 , Eqs. (18) we obtain:

$$\begin{aligned} z_{,\tau}^2 &= \phi^4(zK_2 + 2lK_1)^2 + \phi^4h(K_4^2 - K_3^2), \\ t_{,\tau}^2 &= s\phi^4\{(K_4^2 - K_3^2) + (zK_2 + 2lK_1)^2/h + \\ &\quad (K_2 - K_1M)^2/h\} + \phi^4K_1^2 - \phi^2s\delta. \end{aligned} \quad (39)$$




FIG. 3: It is shown the plot for the metric function $s(t)$ and its derivatives, $\dot{s}(t)$, $\ddot{s}(t)$ for the case $l = 0$. The three functions are bounded at $t = 0$, but $\ddot{s}(t)$ diverges at $at \rightarrow 1$. In this graphic the values of the constants are $A = 0.8$, $C_1 = 0.5$, $b = 0.8$ and $\epsilon = 1$. For $\epsilon = 0, -1$ the behavior changes quantitatively.

For $\epsilon = 0$ we have from the motion constants that imply the Killing vectors in Eqs. (17),

$$\begin{aligned} z_{,\tau}^2 &= 4l\phi^4 h^2 \{lK_3^2 - K_4 K_1 - K_1(lK_1 z^2 - K_2 z)\}, \\ t_{,\tau}^2 &= s\phi^4 \{4l^2 h(K_3 - K_1 y)^2 + (K_2 - K_1 M)^2/h\} + \\ &\quad \phi^4 K_1^2 - \phi^2 s\delta. \end{aligned} \quad (40)$$

In the three cases, $\epsilon = 1, -1, 0$ the geodesics $t(\tau)$ and $z(\tau)$ are complete.

If one considers the topology RXS^3 for Bianchi G_3IX (case $\epsilon = 1$) then $0 < x < 4\pi$ and x reaches a finite value as τ increases. This geodesic “wraps” around the x -direction an infinite number of times. It corresponds to the “phenomena” known as imprisoned incompleteness, described by Hawking (see paragraph 8.5 in [1]) for Taub-NUT spaces. This pathological behavior arises essentially because x is identified at 0 and 4π , a consequence of the compactness of Bianchi type IX.

However both types IX and VIII can be considered non-compact and we can take the range of x being $(-\infty, \infty)$. Being x no longer periodic the geodesics $x(\tau)$ can be extended indefinitely and the spaces are complete. The space VIII resembles the ones studied by Siklos [24] as extensions of Taub-NUT metrics.

B. Kantowski-Sachs and Bianchi spaces G_3I and G_3III

Let us address the case $l = 0$. The included spaces are Kantowski-Sachs and Bianchi spaces G_3I and G_3III for $\epsilon = 1, 0, -1$, respectively.

In this case $\phi = At$ is only a scaling of time coordinate; $s(t)$, Eq.(12), can be integrated in exact form as

$$\begin{aligned} s(t) &= \frac{2\sqrt{2}}{3} Ab^2 t^3 \{K[\frac{1}{2}] - F[\frac{\pi}{2} - \arcsin At, \frac{1}{2}]\} + \\ &\quad \frac{2b^2}{3A^2} (\sqrt{1 - A^4 t^4} - 1) + C_1 A^4 t^3 - \epsilon t^2, \end{aligned} \quad (41)$$

where F is the elliptic integral of the first kind and K is the complete elliptic integral of the first kind. We observe from Eq. (41) that $s(t)$ take real values in the interval $0 \leq (At)^4 < 1$ and that $s(t=0) = 0$.

The metric function $s(t)$, and its derivatives, $\dot{s}(t)$ and $\ddot{s}(t)$ are finite at $t = 0$, then the Weyl scalar Ψ_2 , Eq. (28), and the invariants derived from it are regular at $t = 0$. This is an indication that singularity at $t = 0$ is not a physical one. However, $\ddot{s}(t)$ diverges as $At \rightarrow 1$ and it implies that the polynomial invariants constructed from Ψ_2 diverge as well at $At = 1$, having there the possibility of a singularity. We shall analyze the geodesics to figure out the singularity structure if it exists. Fig. 3 displays the metric function $s(t)$, and its derivatives, $\dot{s}(t)$ and $\ddot{s}(t)$.

The equations for the first derivatives of the coordinates with respect to the affine parameter τ are equally valid in this case, just putting $l = 0$, $M = 0$ and the metric function $s(t)$ from Eq. (41). If the first derivatives are bounded it implies that geodesics are complete and the field is nonsingular [23]; the first equations being,

$$\begin{aligned} y_{,\tau} &= \frac{K_2 \phi^2}{h}, \\ x_{,\tau} &= \frac{K_1 \phi^2}{s}, \end{aligned} \quad (42)$$

The term $\frac{\phi^2}{s}$ that might diverge, is bounded in all the range of t and diverges at $t = 0$ only when $\epsilon = 0$, because

$$\lim_{t \rightarrow 0} \left(\frac{s}{\phi^2} \right) = -\frac{\epsilon}{A^2}, \quad (43)$$

in such a manner that geodesics $x(\tau)$ are unbounded for Bianchi G_3I ($\epsilon = 0$), since $x_{,\tau}$ diverges for a finite value of the affine parameter. Even for the case $\epsilon = 0$ if the constant $K_1 = 0$ then the geodesics $x(\tau)$ are complete since in this case $x(\tau) = A\tau + B$, A, B constants, and it is a complete curve.

The coefficients in equations for $z_{,\tau}$ and $t_{,\tau}$ are bounded as can be seen from expressions (36)-(40) putting $l = 0$, $M = 0$. On the other side, in spite that

$$\lim_{at \rightarrow 1} (\ddot{s}) \rightarrow \infty, \quad (44)$$

all the coefficients in the equations for $x_{,\tau}^\alpha, x^\alpha = (x, y, t, z)$ are bounded, in particular the term $\frac{\phi^2}{s}$ is finite there,

$$\lim_{at \rightarrow 1} \left(\frac{s}{\phi^2} \right) = C_1 A + \frac{2\sqrt{2}b^2}{3A^2} K[\frac{1}{2}] - \frac{2b^2}{3A^2} - \frac{\epsilon}{A^2}. \quad (45)$$

Then at ($at \rightarrow 1$) the geodesics are complete, showing that there is no singularity at that time.

In relation to the electromagnetic fields, it can be seen from Eqs. (14) that when $l = 0$ only electric or only magnetic field persists (remind the electromagnetic duality, which is preserved in the BI theory), for instance, $D(t) = \phi(t)^2$, $B = 0$. But note that the fields can not trespass the threshold of the maximum field b , then $D(t) = A^2 t^2 < b$ imposes an additional restriction for the time coordinate, it must be that $At < \text{Inf} \{\sqrt{b}, 1\}$.

V. SINGULARITY-FREE SOLUTIONS WITH LINEAR ELECTROMAGNETIC FIELD

The conserved quantities derived from the existence of the four Killing vectors are independent of the linear or nonlinear electrodynamics. Then the equations for the derivatives of the coordinates with respect to the affine parameter can be analyzed to test completeness if instead of having the BI field, we take the linear limit.

Expressions for the derivatives of the coordinates with respect to the affine parameter, Eqs. (33)-(40) are valid now with $s(t)$ given by its linear limit, Eq. (29). The possibility of divergence appears again in the term s/ϕ^2 in the equation for $x_{,\tau}$. We analyze the behavior of the term s/ϕ^2 . Integrating the expression for $s(t)$ Eq. (29) with $\phi = A \cos t$ we obtain:

$$s(t) = C_1 \phi^3 \dot{\phi} + \frac{C^2}{A^6} \left[\phi^4 - \frac{5}{3} \phi^2 \dot{\phi}^2 - \frac{A^2}{3} \dot{\phi}^2 \right] + \frac{\epsilon}{A^4} [\phi^4 - \phi^2 \dot{\phi}^2], \quad (46)$$

where $C^2 = H^2 + D^2$ is proportional to the energy density of the field and the limits at $t \rightarrow 0$ and $t \rightarrow \pi/2$ are

$$\begin{aligned} \lim_{t \rightarrow 0} s(t) &= C^2/A^2 + \epsilon, \\ \lim_{t \rightarrow \pi/2} s(t) &= -C^2/3A^2, \end{aligned} \quad (47)$$

From the previous expressions it is clear that $s(t)$ is not zero neither at $t = 0$ nor $t = \pi/2$, being the presence of the electromagnetic field the term that allows the completeness of the geodesics; therefore the coefficients in the equation for $x_{,\tau}$ are bounded and consequently the geodesic $x(\tau)$ is complete. This proves that for the solution of the Einstein-Maxwell equations given by the line element (10) with $s(t)$ in Eq. (46) there are singularity-free spacetimes.

Moreover, for the case $l = 0$ we also have singularity-free solutions since the term s/ϕ^2 is not zero neither at $t = 0$ not at $At = 1$; in this case $\phi = At$ and

$$s(t) = C_1 A^4 t^3 - C^2 - \epsilon t^2, \quad (48)$$

and the limits

$$\begin{aligned} \lim_{t \rightarrow 0} s(t) &= -C^2, \\ \lim_{at \rightarrow 1} s(t) &= C_1 A - C^2 - \epsilon/A^2, \end{aligned} \quad (49)$$

Showing that the geodesics are complete, since the coefficients in the equation for $x_{,\tau}$ are bounded. Summarizing, there are singularity-free solutions with constant electromagnetic field for spacetimes given by the metric (10) and the metric function $s(t)$ given by (46) if $l \neq 0$ or by (48) if $l = 0$. The electromagnetic field is given by Eqs. (30). We also note that for these singularity-free solutions the electromagnetic field tensor does not violate SEC.

VI. CONCLUSIONS

We investigated spacetimes with a four-dimensional group of isometries with a three-dimensional subgroup acting transitively on spacelike hypersurfaces, i. e. spatially homogeneous spacetimes anisotropic in one spatial direction. We address the study of geodesic completeness for solutions to nonlinear electromagnetic field coupled with Einstein equations for cases of Bianchi G_3I , G_3II , G_3III , G_3VIII , G_3IX and Kantowski-Sachs.

Two families are distinguished: Bianchi G_3I , G_3III and Kantowski-Sachs on one side and Bianchi G_3II , G_3VIII and G_3IX on the other.

Bianchi G_3I , G_3III and Kantowski-Sachs only admit one component of the electromagnetic field, electric or magnetic (not both, interchanging roles by a duality rotation). These spaces show divergence in their invariants at a finite time, however, geodesics are complete. At $t = 0$ it is shown that geodesic incompleteness appears for the case $\epsilon = 0$.

Types VIII and IX are geodesically complete and as such they are singularity-free. The space VIII resembles the ones studied by Siklos [24] as extensions of Taub-NUT metrics. When Bianchi G_3IX is considered as having the topology RXS^3 it presents imprisoned incompleteness, described by Hawking (see paragraph 8.5 in [1]) for Taub-NUT spaces. This pathological behavior arises essentially as a consequence of assuming compactness for Bianchi type IX.

Our results are not in contradiction with the assertion by Hawking about the necessity of singularity in non-empty spatially homogeneous models, since the timelike convergence condition fails to hold in nonlinear electrodynamics. The incompleteness due to the presence of imprisoned geodesics in Bianchi G_3IX seems to be inconsistent with the affirmation that: *in the Taub-NUT case the incompleteness of geodesics leads to a real singularity once any amount of matter is added* (see for instance [20]). In this paper we have presented a counterexample for a space of Taub-NUT type with imprisoned geodesics but there is no contradiction since we have considered as

matter a nonlinear electromagnetic field and the original assertion was made thinking of matter as a perfect fluid.

Finally we remark that in the linear limit (Einstein-Maxwell) the solutions are singularity-free for all the Bianchi cases included, being clearly in this case the presence of the field that makes the geodesics complete. In the nonlinear case the determinant factor to avoid the singularity is the curvature parameter, existing singular

solutions if the curvature parameter is zero ($\epsilon = 0$).

Acknowledgments

R. G-S. acknowledges partial support from project UAEM 1941/2004.

-
- [1] Hawking S and Ellis G 1973 *The large scale structure of space-time*, (Cambridge University Press, Cambridge)
 - [2] Geroch R 1968 *Ann. Phys.* **48**, 526.
 - [3] Mars M and Senovilla J M M 1997 *Class. Quantum Grav.* **14** 205.
 - [4] Berger B K 2000 *Phys. Rev. D* **61** 023508.
 - [5] Misner C W 1969 *Phys. Rev. Lett.* **22** 1071.
 - [6] Toporensky A V and Ustiansky V O Dynamics of Bianchi IX universe with massive scalar field: *Preprint* gr-qc/9907047.
 - [7] García-Salcedo R and Bretón N 2000 *Int. J. Mod. Phys. A* **15** 4341.
 - [8] García-Salcedo R and Bretón N 2003 *Class. Quantum Grav.* **20** 5425.
 - [9] Jackson J D 1975 *Classical Electrodynamics* Second Ed. (John Wiley and Sons).
 - [10] Heisenberg W and Euler H 1936 *Z. Phys.* **98** 714.
 - [11] Drummond I T and Hathrell S J 1980 *Phys. Rev. D* **22** 343.
 - [12] Schwinger J 1951 *Phys. Rev.* **82**, 664.
 - [13] A. A. Tseytlin A A 1996 *Nucl. Phys. B* **469** 51; Fradkin E S and Tseytlin A A 1985 *Phys. Lett. B* **163** 123.
 - [14] Brill D R 1964 *Phys. Rev.* **133** 845.
 - [15] Hughston L and Jacobs K 1970 *Astrophys. J.* **160** 147.
 - [16] Hacyan S 1984 *Astron. Astrophys.* **140** 363.
 - [17] Plebański J F 1970 *Lectures on Nonlinear Electrodynamics* (Copenhagen).
 - [18] Plebański J F 1973 *Spinors, Tetrads and Forms* (Ed. CINVESTAV-IPN, México).
 - [19] Stephani H, Kramer D, MacCallum M, Hoenselaers C, Herlt E, 2003 *Exact Solutions of Einstein's Field Equations*. (Cambridge University Press).
 - [20] Ryan M P and Shepley L C 1975 *Homogeneous Relativistic Cosmologies* (Princeton Univ. Press, Princeton).
 - [21] Senovilla J M M 1988 *Gen. Rel. and Grav.* **30** 5701.
 - [22] Visser M and Barceló C 2000 Energy conditions and their cosmological implications *Preprint* gr-qc/0001099.
 - [23] Arnold V I 1990 *Ordinary Differential Equations*, (MIT Press).
 - [24] Siklos S T C 1976 *Phys. Lett. A* **59** 173.